

Lembulus forlii n. sp. from the Pliocene of Italy (Bivalvia, Nuculanidae)

Eivind Palm* & Bruno Dell'Angelo[#] (✉)

* Byvej 16, 4591
Føllenslev, Denmark
eipa@km.dk

Via Santelia 55/12A,
16153 Genova, Italy
bruno.dellangelo@
chitons.it
(✉) Corresponding
Author

Abstract

A new *Lembulus* species (Bivalvia, Nuculanidae) is described from the Pliocene (Zanclean-Piacenzian) of Northern Italy. *Lembulus forlii* n. sp. is closely similar to *Lembulus pella* (Linné, 1758), from which it differs mainly by being more elongate, with a longer, curved rostrum and a finer sculpture.

Riassunto

Viene descritta una nuova specie di *Lembulus* (Bivalvia, Nuculanidae) del Pliocene (Zancleano-Piacenziano) italiano. Il materiale proviene da tre località, due in Toscana (Spicchio e Cedda) ed una in Emilia Romagna (Lugagnano). *Lembulus forlii* n. sp. è notevolmente simile a *Lembulus pella* (Linné, 1758), l'unica specie in precedenza nota per questo genere nel Plio-Pleistocene. La nuova specie si distingue per la conchiglia più allungata, il rostro più lungo e curvo e la scultura più fine.

Key words

Bivalvia, Nuculanidae, *Lembulus*, new species, Pliocene, Italy.

Introduction

In recent years some investigations on the protobranch of the Mediterranean Sea have been carried out (e.g. Di Geronimo & La Perna, 1996; La Perna, 2003, 2008; Della Bella & Tabanelli, 2004; La Perna et al., 2004). These papers mainly deal with Pleistocene to Recent deep water species, and only one of them (La Perna, 2008) includes the genus *Lembulus*, without new species described. For some years, one of the authors (E.P.) has been collecting a large amount of material from several Italian Pliocene localities and in the process of sorting out the species, some new species have been encountered. Among them, there is a distinct member of the genus *Lembulus*, here described as a new species. The genus *Lembulus*, described by Risso (1826), is characterized by having a robust, strongly convex shell, with a short, stout, bicarinate rostrum terminating into a pointed tip. The commarginal sculpture is weak, but a dense pattern of well defined, oblique striae is present. It ranges from the Early Miocene (Cossmann & Peyrot, 1913) and is still present in the Atlantic (from Portugal to West Africa, south to Angola), including the Mediterranean (Poppe & Goto, 1993; Giannuzzi-Savelli et al., 2001; La Perna, 2008). The type was fixed by Gray (1847) as *Lembulus rossianus* Risso, 1826, a synonym of *Arca pella* Linné, 1758 (Arnaud, 1977; La Perna, 2008). *Lembulus pella* was the only species known for the genus in Plio-Pleistocene of the Mediterranean area. *Arca interrupta* Poli, 1795, *Nucula rostrata* Borson, 1823 and *L. pella* var. *anterotunda* Sacco, 1898 are all synonyms of *L. pella* (La Perna, 2008).

Tuscany (Spicchio and Cedda) and one in Emilia Romagna (Lugagnano). Spicchio (Firenze) is an outcrop near Empoli, of Piacenzian age (Dominici et al., 1997). Cedda (Siena) is an outcrop near Poggibonsi (5 km East), referred to the Early-Middle Pliocene (Brunetti & Della Bella, 2008). Lugagnano (Piacenza) is a classic Pliocene locality near Piacenza, of Zanclean age (Rio et al., 1988).

The following abbreviations are used: BD = Bruno Dell'Angelo collection (Genova, Italy); EP = Eivind Palm collection (Denmark); IGF = Museo di Storia Naturale, Sezione di Geologia e Paleontologia, Firenze (Italy); MZB = Museo di Zoologia dell'Università di Bologna, Bologna (Italy).

Dimensions (in mm) are given as length × height × width (for complete shells) or length × height (for single valves).

Systematics

Class Bivalvia Linné, 1758
Subclass Protobranchia Pelseneer, 1889
Order Nuculoida Dall, 1889
Family Nuculanidae H. & A. Adams, 1858
Genus *Lembulus* Risso, 1826 ex Leach ms
Type species: *Arca pella* Linné, 1758

Lembulus forlii n.sp.
(Fig. 1A-H)

Type material

Holotype: Spicchio (Piacenzian), 1 valve (MZB 45686, 11.3 × 5.7 mm). Paratypes: Spicchio (Piacenzian), 2 shells

Material and methods

The material originated from three localities, two in

(max $13 \times 7 \times 2.7$ mm), 28 valves (EP, max 13×7 mm), 2 valves (MZF 45687, respectively 10.8×5.5 and 8.3×4.4 mm), 1 valve (BD, 9.8×5 mm); Lugagnano (Zanclean), 1 shell (IGF 15692E, $8 \times 4.3 \times 3.5$ mm), 2 shells (max $10 \times 5 \times 3.5$ mm), 1 valve (EP, 11×6 mm); Cetona (Pliocene), 21 valves (EP, max 12.2×6.5 mm).

Description

Medium sized, inaequilateral, aequivalve, rather robust, moderately convex shell. Umbo opisthogyrate, at about 2/5 from posterior margin. Rostrum long, slightly bent, making the length about twice the height; rostrum tip somewhat pointed. A rounded, rather strong keel bounds the rostrum dorsally; another weaker keel from umbo to postero-ventral margin, defines a slightly concave rostral area. Anteriorly, a weak radial keel. Ornamentation oblique, consisting of closely spaced ribs; growth striae irregularly spaced. Resilifer small, triangular. Hinge plate moderately thick. Anterior teeth row a little shorter than posterior. Posterior row reach about 2/3 of the total posterior margin. Hinge strong with chevron-shaped

teeth, 14 anteriorly, 18 posteriorly in the holotype. Pallial line not well defined; pallial sinus rounded, wide, rather deep. Muscle scars well impressed.

Type locality

Spicchio (Firenze, Tuscany).

Distribution

Lembulus forlili n.sp. is so far only known from the three localities mentioned above, all of Pliocene age (Zanclean-Piacenzian).

Etymology

Named after our good friend Maurizio Forli.

Discussion

Lembulus forlili n.sp. (Fig. 1A-H) is notably similar to *L. pella* (Fig. 1K, L), but the shell is more elongate, and the rostrum markedly longer and more pointed in the new species. The umbo is clearly closer to the center in *L.*




Fig. 1. A-H. *Lembulus forlili* n. sp. **A, B.** Spicchio (Firenze), length 7 mm; **C, D.** Holotype (MZF 45686), Spicchio (Firenze), length 11.3 mm; **E, F.** Spicchio (Firenze), length 6.2 mm; **G, H.** Cetona (Siena), length 11 mm; **I, J.** Paratype (IGF 15692E), Lugagnano (Piacenza), length 8 mm (both valves); **K, L.** *Lembulus pella* (Linné, 1857), Pietrafitta (Siena), length 11 mm.

Fig. 1. A-H. *Lembulus forlili* n. sp. **A, B.** Spicchio (Firenze), lunghezza 7 mm; **C, D.** Olotipo (MZF 45686), Spicchio (Firenze), lunghezza 11.3 mm; **E, F.** Spicchio (Firenze), lunghezza 6.2 mm; **G, H.** Cetona (Siena), lunghezza 11 mm; **I, J.** Paratipo (IGF 15692E), Lugagnano (Piacenza), lunghezza 8 mm (entrambe le valvole); **K, L.** *Lembulus pella* (Linné, 1857), Pietrafitta (Siena), lunghezza 11 mm.

pella, i.e. the shell is distinctly more inaequilateral in the new species. Even in younger specimens (Fig. 1E, F), the rostrum is longer and the umbo is more posterior than in *L. pella*. The keel impression is weaker, and the posterior bicarinate shape typically seen in *L. pella*, is very weak or absent in the new species. The ornamentation in *L. forlpii* n.sp. is tighter than usual in *L. pella*, though *L. pella* is very variable in that respect. The whole surface gives an impression of smoothness, whereas in *L. pella*, the ribs are slightly more pronounced. In *L. forlpii* n.sp., the hinge rows make a wider angle and the ligament pit is notably smaller, than in *L. pella*.

Lembulus emarginatus (Lamarck, 1819) perhaps belong to yet another, may be a Miocene species, but it is unclear what Lamarck really meant, when he described the species (for a more detailed discussion, see La Perna, 2008). The figures in literature, attached to that name, are very similar to *L. pella* and they do not have the long, bent rostrum, so characteristic in *L. forlpii* n.sp.

Lembulus undatus (Defrance, 1825), a Miocene species, has coarse, transversal ridges, giving the appearance of a wavy surface. Some specimens we have seen (from Aquitaine, South of France) have more, other less of these ridges, but *L. pella* and *L. forlpii* n.sp. never has such a sculpture.

Lembulus forlpii n. sp. is only found in clayey deposits, whereas we have seen thousands specimens of *L. pella* and none of the new species in the sandy deposits in the Siena-Firenze area. The presence of *L. pella* is well documented in the Italian Pliocene and Pleistocene (Sacco, 1898; Cavallo & Repetto, 1992; La Perna, 2008). At Spicchio, there are some sandy-clayey lenses where *L. pella* can occasionally be found. These shells show the typical characters of this species, whereas neither *L. forlpii* n.sp. nor intermediate forms have been found. These observations support the interpretation of *L. forlpii* n.sp. as a distinct species, and not as an ecophenotype of *L. pella* which lived in muddy bottoms or deeper waters. Also at Melograni (near Pietrafitta and Bibbiano, San Gimignano), where the sediment is richer in pelitic fraction than in most of the other sandy localities in the Siena area, typical shells of *L. pella* are found, whereas *L. forlpii* n.sp. has been never found.

The two valves we have seen from Ciuciano, a place with deeper water deposits (Forli & Dell'Angelo, 2000), are *L. pella*, but the species is very rare here, presumably because it was too deep for optimal living conditions. Most probably, due to its close resemblance to *L. pella*, *L. forlpii* n.sp. has been always confused with the former and its stratigraphic distribution could be wider than that hitherto known.

Acknowledgements

We wish to thank Antonio Bonfitto (MZB) who prepared the illustration and Birgitte Palm, wife of the first author, for the photographs. We especially thank Rafael La Perna (Università di Bari) for his helpful comments

and constructive criticism, that greatly improved the quality and completeness of the manuscript.

References

- ARNAUD P.M., 1977. Révision des traxa malacologiques Méditerranéens introduit par Antoine Risso. *Annales du Muséum d'Histoire Naturelle de Nice*, **5**: 101-150.
- BRUNETTI M. & DELLA BELLA G., 2008. *Macalia (?) kengii* n. sp. un nuovo bivalve per il Pliocene toscano (Tellinidae, Macominae). *Bollettino Malacologico*, **44**: 115-118.
- CAVALLO O. & REPETTO G., 1992. Conchiglie fossili del Roero. Atlante Iconografico. *Memorie dell'Associazione Naturalistica Piemontese*, **2**: 1-251.
- COSSMANN M. & PEYROT A., 1913. Conchologie Néogénique de l'Aquitaine. *Actes de la Société Linnéenne de Bordeaux*, **66**: 121-324.
- DELLA BELLA G. & TABANELLI C., 2004. Revisione della malacoafuna pliocenica di Rio Albonello (Brisighella, Ravenna). III. *Testyleda amaritae* n. sp. *Quaderno di Studi e Notizie di Storia Naturale della Romagna*, **19**: 7-16.
- DI GERONIMO I. & LA Perna R., 1996. *Bathyspinula excisa* (Philippi, 1844) (Bivalvia, Protobranchia): a witness of the Plio-Quaternary history of the deep Mediterranean benthos. *Rivista Italiana di Paleontologia e Stratigrafia*, **102**: 105-118.
- DOMINICI S., MAZZANTI R. & NENCINI C., 1997. Geologia dei dintorni di San Miniato tra l'Arno, l'Elsa e l'Era. *Quaderni del Museo di Storia Naturale di Livorno*, **14** (suppl. 1): 1-27.
- FORLI M. & DELL'ANGELO B., 2000. A new species of *Marginella* (Mollusca, Gastropoda) from the Italian Pliocene. *Bollettino Malacologico*, **36**: 93-98.
- GIANNUZZI SAVELLI R., PUSATERI F., PALMERI A. & EBREO C., 2001. *Atlante delle conchiglie marine mediterranee*. 7. *Bivalvia: Protobranchia-Pteriomorpha*. Edizioni Evolver, Roma: 246 pp.
- GRAY J.E., 1847. A list of the genera of recent Mollusca, their synonymy and types. *Proceedings of the Zoological Society of London*, **15**: 129-206.
- LA Perna R., 2003. The Quaternary deep-sea protobranch fauna from the Mediterranean: composition, depth-related distribution and changes. *Bollettino Malacologico*, **39**: 17-34.
- LA Perna R., 2008. Revision of the Nuculanidae (Bivalvia: Protobranchia) from the Cerulli Irelli collection (Mediterranean, Pleistocene). *Bollettino della Società Paleontologica italiana*, **46**: 125-137.
- LA Perna R., CEREGATO A. & TABANELLI C., 2004. Mediterranean Pliocene protobranchs: the genera *Jupiteria* Bellardi, 1875, *Ledella* Verrill & Bush, 1897 and *Zealeda* Marwick, 1924 (Mollusca, Bivalvia). *Bollettino Malacologico*, **40**: 25-36.
- POPPE G.T. & GOTO Y., 1993. *European Seashells. Volume II (Scaphopoda, Bivalvia, Cephalopoda)*. Christa Hemmen Verlag: 221 pp.
- RISSO A., 1826. *Histoire naturelle des principales productions de l'Europe méridionale et principalement de celles des environs de Nice et des Alpes-Maritime. Mollusques*. E.G. Levrault, Paris: 439 pp.
- RIO D., SPROVIERI R., RAFFI S. & VALLERI G., 1988. Biostratigrafia e paleoecologia della sezione stratotipica del Piacenziano. *Bollettino della Società Paleontologica Italiana*, **27**: 213-238.
- SACCO F., 1898. *I molluschi dei terreni terziarii del Piemonte e della Liguria. Parte XXVI*. Carlo Clausen Torino: 93 pp.