



An area-analytical zoogeographical classification of Palearctic Unionaceae species

Károly Bába

KEY WORDS: Unionaceae, area-analytical division, fauna circles, West-Palearctic-, East-Palearctic-Elements, Pacific-Palearctic-Elements.

ABSTRACT: The zoogeographical classification for 31 *Unionaceae* species is given by using the area-analytical zoogeographical approach formerly successfully applied to terrestrial snails (Bába 1982). Studied species belong to the following genera: *Margaritifera*, *Margaritana*, *Unio*, *Potamida*, *Lanceolaria*, *Pseudanodonta*, *Anodonta*, *Cristaria* and *Corbicula*.

RIASSUNTO: La classificazione zoogeografica di 31 specie di unionacea è stata stilata utilizzando un metodo area-analitico. Le specie appartengono agli zonobiomi Boreale e Aboreale. Nei due maggiori gruppi dello zonobiotma Aboreale - i.e. nei gruppi Paleartici dell'Ovest e dell'Est- gli elementi che si originano dalla Siberia orientale, centrale e occidentale, circa 20 specie, sono i più abbondanti. Sono necessarie ulteriori ricerche per determinare l'origine degli elementi della Manciuria e Sinopacifici nei gruppi della Siberia orientale (15 specie). I gruppi occidentali paleartici includono gli elementi centro-asiatici, ponto-caspici e olomediterranei, le specie Adriatico e Atlanto Mediterranea da rifugi secondari. La maggior parte di queste specie posseggono sia areali stabili che in regressione.

KÁROLY BÁBA, Gyula Juha'sz teachers training college, Dep. of biology Vár u. 6., Szeged, Hungary.

INTRODUCTION

In contrast to numerous invertebrate taxa (*Plecoptera*, *Heteroptera*, *Odonata*, *Ephemeroptera*, *Trichoptera*, etc.), the area-analytical classification of freshwater molluscs has not been prepared yet. This gap is filled partly in this study for the *Unionaceae*, and was done so earlier for *Sphaeriidae* (Bába 1997).

MATERIAL AND METHODS

Earlier studies proved that the dispersal of terrestrial and aquatic organisms follow the same fundamental rules (De Lattin 1967, Varga 1975, Dévai 1976), that also holds true for plants and terrestrial snails (Bába 1986). The zoogeographical categorisation of species is further hampered by the limited information available on the subspecies distinguished by mostly anatomical traits. Albeit, the geographical distribution of subspecies carries important information on the history of the species (Heptner 1959). Different subspecies reported from the same area in the literature are sympatric, that in turn makes the subspecies status questionable. The dispersal of clams was enhanced by certain human activities, like fish introduction (*A. woodiana*) and waterway canalizations (*Dreissena polymorpha*). Occasionally, the glochidium may be transported by migrating birds. Nevertheless, the fossil-based dynamics for East Palearctic species is incomplete, which could serve with essential information on their distribution.

Generalised maps for the species (Fig. 2-5) were prepared by using literature data from the listed references. Fauna elements and refugium areas (Fig. 1) were classified after Dévai (1976). Due to the occurrences of various combinations, Manchurian refugia in East Siberia were collectively considered as East Siberian elements. For faunistic data from Spain, Portugal and Switzerland I thank F. Ramires and H. Turner. The help of S. Bagdi in drawing the maps is also acknowledged.

FAUNA GROUPS

The 31 studied *Unionaceae* species belong to the Boreal and Aboreal zonobiomes. The range of the Boreo-Alpine circumpolar (Holarctic) *Margaritifera margaritifera* from the Boreal zonobiome (Fig. 2) was drawn based on the work of Jungbluth et al. (1985). In the Aboreal zonobiome the most species rich fauna element group in the East Palaearctis is the East Siberian, which includes many refugia. These refugia may combine with each other. The Amurian (Manchurian) elements are of particular importance, since these extended from the Amurian region into North China, Japan and Korea through the Japan Inner Lake during the Mindel Glaciation (De Lattin 1967). This resulted in the Manchurian-Sinopacific distribution pattern, which is temporarily classified as East Siberian elements based on De Lattin's theory.

Shadin (1952) gives no information on the subspecies. Species assigned to this category are *Margaritana middendorfi* Rosen 1926, *Margaritana sacchariensis* Shadin 1938, *Margaritana daburica* (Middendorf 1850), *Unio continentalis* (Haas 1980), *Lanceolaria grayana* Lea 1834, *Lanceolaria cylindrica* (Simpson 1900), *Anodonta beringiana* Middendorf 1850, *Anodonta encaphys* Hende 1879, and *Anodonta arafurica* Hende 1877. These fauna elements have stationary areas (Figs 2-3). Species with Manchurian-Sinopacific disjunct area are *Unio douglasiae* Griffit at Pitaeon 1833, *Anodonta woodiana* (Lea 1834) (adventives in Europe arrived with Amurian fish), *Cristaria plicata* (Leach 1815), *Corbicula fluminea* (O.F. Müller 1774), Fig. 3.

Anodonta anatina (Linné 1758) is a Middle Siberian element from the Angarian refugium (Fig. 3). West Siberian elements most probably originated in the West Siberian meltwater lake, and then distributed toward east and - through a link with the Onega Lake - in the direction of Europe during the Riss glaciation (De Lattin 1967). Such species are : *Unio tumidus* Retzius

West-Palearctic Elements

1. South-mediterranean Elements
(Canarian, Mauritanian, Tyrrhenian, Cyprean, Cyprian Refugial areas)
1. Holomediterranean Elements
 - 1.a Atlantomediterranean
 - 1.b Adriatomediterranean Refugial areas
 - 1.c Pontomediterranean
 - 1.d South Italian
 - 1.e euxin
2. Ponto-Caspian Elements
Ponto-Csapian Refugial area
- West-Asian Elements
 - a) Pre-Asian Elements
3. Syrian Refugial areas
4. Iranian
- b) Central-Asian Elements
5. Afghan Refugial areas
6. Turkestanian

East-Palearctic Elements

9. Mongolian Elements
Dzsungarian Refugial areas
Mongolian-Altaic-Hangayn Refugial areas
10. Siberian Elements
 - a) West Siberian Elements
 7. West Siberian Refugial area
 - b) Central Siberian Elements
 8. Angaran Refugial area
 - c) East Siberian Elements
 10. c) East Siberian Elements
Stanovoy-Bureyan
Okhostkian Refugial areas
Kamchatkan
11. d) Manchurian Elements
Amurean
Sakhalin-Kurilian
Hokkaidon Refugial areas
Manchu-Ussurian

Pacific-Palearctic Elements

12. Japanese
13. Korean
14. Sino-Pacific Refugial areas
15. Sino-Tibethian
16. Yunnan

Note: The Korean Refugial area belongs to Manchurian elements by De Lattin 1967.

Fig. 1 System of freshwater refugial areas (fauna circles) and fauna elements in the Arboreal of Palearctic region (DE LATTIN 1967, Z. VARGA 1975) from GY. DÉVAI 1976.



1788, *Unio crassus* Retzius 1788, *Unio pictorum* (Linné 1758) and *Anodonta cygnea* (Linné 1758). The subspecies appear at the edges of the species' European ranges in Western and Central Europe and in the Pontis, arranged radially with the former Onega Lake in center (Figs 3-4).

West Palearctic elements include the Central Asian (Afghan) fauna group element *Corbicula fluminalis* (O.F. Müller 1774), which colonised Central Europe during the Pleistocene (Mindel-Riss interglacial) already, since its fossils have been found in Germany, Czech Republic and Hungary (Lozek 1964, Zilch et al. 1960). The species' recent range (Fig 4) is given by Girardi (1989-90). Central Asian elements originate from non-European territories of the great Mediterranean refugium (De Lattin 1967).

Stationary species in the Ponto-Kaspian group: *Unio stevenianus* Kyrnicky 1837, *Unio sieversi* Drouët 1881, *Unio mingrelicus* Drouët 1881, *Anodonta cyrea* Drouët 1881 and the disjunct *Pseudanodonta complanata* (Rossmässler 1835). (Fig 4)

For Holomediterranen elements the following division can be made. Adriato-Mediterranen refugium: *Microcondylea compressa* (Menke 1836). Atlanto-Mediterranean refugium: *Margaritifera auricularis* (Sprengler 1793). Great Holomediterranean refugium: *Unio mancus* Lamarck 1819, *euxin Unio hueti* (Bourguignat 1836) and the regressing *Potamida littoralis* (Lamarck 1801). (Figs 4-5).

SUMMARY

The zoogeographical classification for 31 *Unionaceae* species was completed using the area-analytical method. The species belong to the Boreal and Aboreal zonobiomes. In the two major fauna groups within the Aboreal zonobiose - i.e. in the West and East Palearctic groups - fauna elements originating from East-, Middle- and West Siberian refugia are the most abundant, altogether 20 species. Further research is necessary to determine the origin of Manchurian Sinopacific elements in the East Siberian group (15 species).

The West Palearctic group includes Central Asian, Ponto-Kaspian and Holomediterranean elements, and Adriato- and Atlanto-Mediterranen species from the secondary refugia. Most of these species posses either stationary or regressive area.

LITERATURE

- ADAM, W. (1960): Mollusques terrestres et dulcicoles. *Faune de Belgique I.* Brüsselles, 1-402.
- AKRIMOVSKIJ, N. N. (1976): Fauna Armjanskoy CCR (Mollusca). *Akad. Nauk Armjanskoy CCR.* Jerevan, 1-378.
- ALVAREZ, J. - SELGA, D. (1967): Observaciones sobre intertebrados dulceacuicolas de las alrededores de Madrid. *Bol. R. Soc. Espanola Hist. Nat. (Biol.)*, 65, 171-197.
- AMONI, D. - BARLETTA, G. - BIANCHI, J. - BONA, E. - GIROD, A. - MARIANI, M. - TORCHIO, M. (1978): La Malacofauna di Alcuni Loghi insubrici Minori. *Natura Bresciana. Ann. Mus. Civ. St. Nat. Brescia*, 15, 95-119.
- ANT, H. - JUNGBLUTH, J. H. (1979): E.I.S. Beiträge aus Bundesrepublik Deutschland, *Malacologia* 18, 185-195.
- BÁBA, K. 1982. Eine neue Zoogeographische Gruppierung der Ungarischen Landmollusken und die Wertung des Faunabildes. *Malacologia* 22(1-2), 441-454.
- BÁBA, K. (1997): An area-analytical zoogeographical classification of Bivalves in the Sphaeriidae family. Abstract band von Symposium "Ökologie und Taxonomie von Süßwassermollusken" International Congress on Palearctic Mollusca, Salzburg, 4.
- BELLAVERE, C. - PERETTI, E. (1984): Revisione, catalogo e distribuzione al 1878 dei Molluschi viventi nel Versante settentrionale Dell'Appenino dal Tidone al Sacchia della Collectione Strobel. Museo di storia Naturale, Univ. di Parma, Pavona, 54-107.
- BILGIN, F. (1980): Systematics and distribution of Mollusca species collected from some freshwaters of West Anatolia. *Diyarbakir Univ. Tip. Fakultesi Dergesi*, Diyarbakir, 8(2), 1-64.
- BOETTGER, C.R. (1954): Süßwassermuscheln von der Insel Sylt (Nordfrizische Inseln). *Arch. Moll. Frankfurt a Main*, 83, 4-6, 139.
- BOLE, J. (1962): Die Mollusken des Triglav Nationalparks an der Umgebung (Mollusca: Gastropoda, Bivalvia), Varstvonaarave, Ljubljana, 57-58.
- CASTAGNOLO, L. - FRANCHINI, D. - GIUSTI, F. (1980): Bivalvi (Bivalvia) Guide per il Riconoscimento delle specie animali delle acque interne Italiane 10. Consiglio Nazionale delle Ricerche, 1-64.
- COSSIGNANI, V. - COSSIGNANI, T. (1995): Atlante delle Conchiglie terrestri e dulciacquicole Italiane. L'Informatore Piceno, Ancona, 1-208.
- DE LATTIN, G. (1967): Grundiss der Zoogeographie. Gustav-Fischer Verlag, Jena, 1-602.
- DÉVAI GY. (1976): Magyarországi szitakötő (Odonata) fauna chorologai vizsgálata. - The chorological study of the dragonfly (Odonata) fauna of Hungary. *Acta Biol. Debrecina, Debrecen*, 13(1), 119-157.
- DHORA, D. H. - WELTER-SCHULTES, F. W. (1996): List of species and atlas of the non-marine molluscs of Albania. *Schriften zur Malakozoologie, Cismar, Ostholstein* 9, 1-6, 90-197.
- FALCO, G. - CASTAGNOLO, L. (1983): I molluschi viventi, terrestri d'acqua dolce, nello studio biogeographic dell' isola di Sardegna. *Livori della Società Haliana di Biogeographia VIII*, 227-249.
- FLASAR, I. (1991-92): Die Flussperlmuschel Margaritifera margaritifera (L) in dem Ascher Gebiet (A sky vybék) Eulamellibranchiata: Margaritiferidae, *Sbornik* 13-14, 7-25.
- FLASAR, I. (1992): Frühere Verbreitung der Flussperlmuschel Margaritifera Margaritifera (L) im Friedlander gebiet in Nordböhmen (Eulamellibranchiata: Margaritiferidae). *Malakologische Abhandlungen, Dresden* 16 (12), 83-87.
- FORCART, L. (1965): Rezente Land und Süßwassermollusken der süditalienischen Landshaften Apulien, basilicata und Calabrien. *Vervandl. Naturf. Ges. Basel* 76 (1), 59-184.
- FÜKÖH, L. - KROLOPP, E. - SÜMEGI, P. (1995): Quaternary Malakostriatigraphy in Hungary. *Malacological Newsletter*. Gyöngyös. Suppl. 1, 1-219.
- FRANK, C. (1986): Die Molluskenfauna des Kamptales. Studien und Forschungen aus dem Niederösterreichischen Institut für Landeskunde, Wien, 9, 1-118.
- GERMAIN, L. (1931): Faune de France 22. Mollusques terrestres et flu-











Fig. 2 The distribution-maps of the species of the *Margaritifera*, *Margaritana*, *Unio* geni, Boreo-Alpine, Eastern-Siberian (Amurian) fauna elements.








Fig. 3 The distribution-maps of the species of the *Lanceolaria*, *Cristaria*, *Corbicula*, *Unio* geni. East-Siberian (Mandsurian-Sino-Pacific), West-Siberian fauna elements.








Fig. 4 The distribution-maps of the species *Unio*, *Corbicula*, *Pseudanodonta*, *Margaritifera* geni. Middle-Siberian, Ponto-Caspian, Middle-Asian, Adriato-Mediterranean fauna elements.




Fig. 5 The distribution-maps of the species *Unio*, *Potamida* geni. Holomediterranean fauna elements.

- vuiatiles. Federation Francaise des soc. de Sci. naturelles. office central de Faunistique, Paris, pp. 714-778.
- GIRARDI, H. (1989-90): Deux bivalves d'eau douce récents pour la faune française (Mollusca, Bivalva). Bull. Soc. et Sci. nat., Vaucluse, 87-93.
- GIUSTI, F. - CASTAGNOLO, L. (1983): I molluschi viventi terrestri d'acqua dolce, nello studio biografico dell'isola di Sardegna. Lavori della Società Italiana di Biogeografia VIII, 227-249.
- GLÖER, P. - MEIER-BROOK, C. - OSTERMANN, O. (1992): Süsswassermollusken D. J. N. Hamburg, 10, 1-111.
- HEPTNER (1959): Centres of speciation in the fauna of palearctic steppe and desert zone. Proc. XI. Zool. Congr. London 1958, 155-157.
- JAECHEL, S.H. (1955): Die Wassermollusken der Nuthe-Niederung und des Raumes zwischen mittlerer Elbe und Warthe. Abhandlungen und Berichte für Naturkunde und Vorgesichte, Magdeburg IX, 5, 185-217.
- JAECHEL, S. G. - KLEMM, W. - MEISE, W. (1957): Die Land und Süßwassermollusken der nördlichen Balkanhalbinsel. Abhandlung und Berichte aus dem Staatlichen Museum für Tierkunde Forschungstelle. Dresden, 23 (2), 141-205.
- JUNGBLUTH, J. H. - COOMANS, H.E. - GROHS, H. (1985): Bibliographie der Flussperlmuschel Margaritifera margaritifera (Linnaeus 1758) (Mollusca, Pelecypoda) - Verschlagene Technische Gegevens, Institut voor taxonomisch Zoölogia (Zoölogisch Museum). Universiteit van Amsterdam, 41, I-XXXI, 1-220.
- KERNAY, M.P. (1976): Atlas of the non-marine mollusca of the British Isles. Graphic Art., Cambridge, 1-202.
- KLEMM, W. (1960): Mollusca VII a Catalogus faunae Austriae. Akad. Wiss., Wien, 1-59.
- LINDHOLM, A.W. (1901): Beiträge zur Kenntnis der Weichterfauna Südrusslands. Nachrichtenblatt d. deutschen Malakozoologischen Gesellschaft, 11-12, 11-192.
- LINDHOLM, A.W. (1903): Zur Molluskenfauna der Gouvernements Kurks und Orenburg. Ann. du Musée Zoologique 8, 338-344.
- LISICKY, M.J. (1991): Mollusca Slovenska. VEDA Slovenskej Akad. vied., Bratislava, 1-341.
- LOZEK, V. (1964): Quartermollusken der Tschechoslowakei Tschechoslowakischen Akad. der Wiss. Praha, 1-374
- LOZEK, V. (1965): Entwicklung der Molluskenfauna der Slowakei in der Nacheiszeit. Informationsbericht der Landwirtschaftlichen Hochschule Nitra I. 1-4, 9-24.
- MIENIS, H.K. (1986): A revised checklist of the brackish and freshwater Molluscs from Israel and the administered areas, Levantina, 63, 675-682.
- NAGEL, K. - HOFFMEISTER, U. (1986): Microcondylea compressa Menke 1928, Bivalvia: Unionidae recovered alive in Northern Italy. - Bool. Malacologica Milana 22, (9-12), 251-260.
- NESEMANN, H. (1993): Zoogeographia und Taxonomie der Muschel-Gattungen *Unio* Philipsson 1788, *Pseudanodonta* Bourguignat 1877, und *Pseudunio* Haas 1910 im oberen und mittleren Donausystem (Bivalvia: Unionidae, Margaritiferidae). Nachr.



- bl. erste Voralberger Malak. Ges. 1, 20-40.
- PIEHOCKI, A. - DYDUCH-FALINOWSKA, A. (1993): Mieczaki (Mollusca) Male (Bivalvia). Fauna Stodkowodna Polski, Wydawnietwo, Naukowe PWN, Warszawa, 1-204.
- PINTÉR L. - RICHNOVSZKY A. - SZIGETHY A. (1979): A magyarországi recens puhatestűek elterjedése. Soósiana 1, 1-351.
- SÁRKÁNY-KISS, A. (1997): The present-day situation of the Unionidae (Mollusca, Bivalvia) in the Transylvanian tributaries of the river Tisza. Trav. Mus. Hist. Nat. Grigore Antipa 38, (in print)
- SCHLESCH, H. - KNAUSP, C. (1938): Zur Kenntnis der Land- und Süßwassermollusken. Litanens. Arch. moll. Frankfurt a Main. 70 (2-3), 73-160.
- SHADIN, V.C. (1952): Moljuszki presznüh I scolonovatüh bod CCCR. Akademii Nauk CCCR. Moszkva-Leningrad, 1-373.
- SOÓS L. 1943. A Kárpát-medence Mollusca faunája Magyar Tud. Akad. Budapest, 1-478.
- STROUHAL, H.K. - KLEMM, W. (1960): Catalogus Faunae Austriae VIII.a. Mollusca, Springer-Verlag, Wien, 1-60.
- VARGA Z. (1971): A szétterjedési centrumok és a szétterjedési folyamat jelentősége a földrajzi izoláció kialakulása és a mikroevolúció szempontjából. Állattani Közlemények LVIII 1-4, 142-149.
- VARGA, Z. (1975): Geographische Isolation und Subspeziation bei den Hochgebirgslepidopteren der Balkanhalbinsel. Acta Entomol. Jugosl. 11, 1-2, 5-40.
- ZDUN, V.J. (1960): Do fauni Moljuszkiv Zakarpatja. Naukovi Zapuszczi Hankovo-prirodoznavjaoro A. H. YRCR. VIII., 83-94.
- ZILCH, A. (1967): Die Typen und Typoide des Natur-Museum Schenkenberg 39, Mollusca, Unionaceae, Arch. Moll. Frankfurt a Main 97 (1-6), 45-154.
- ZILCH, A. (1983): Die Typen und Typoide des Natur-Museum Schenkenberg, Mollusca: Unionaceae (Nachtrag zu Teil 39), Arch. Moll. Frankfurt a Main 114 (1-3), 77-92.
- ZILCH, A. - Jaeckzel, S. G. A. (1960): Die Tierwelt Mitteleuropas, Mollusken (Ergänzung), Quelle-Meyer, Leipzig, 1-260.